



SURFACE DISINFECTANT

• Fungicide • Mildewstat • Virucide* • Cleaner • Deodorizer

For Home, Hospital, Institutional and Industrial Use

- One-step disinfectant cleaner⁵ is effective against a broad spectrum of bacteria, is virucidal*, fungicidal and inhibits the growth of mold and mildew, when used as directed.
- Use to remove odors caused by flooding and smoke from fires.
- Kill germs, bacteria and viruses* on pre-cleaned environmental surfaces.
- Kills 99.9% of SARS-CoV-2², which causes COVID-19.
- Effective in the presence of 5% serum contamination

ACTIVE INGREDIENTS:

Octyl decyl dimethyl ammonium chloride.....	0.025%
Diocetyl dimethyl ammonium chloride.....	0.010%
Didecyl dimethyl ammonium chloride.....	0.015%
Alkyl (C14 50%; C12 40%; C16 10%)	
Dimethyl benzyl ammonium chloride.....	0.034%
Other Ingredients	99.916%
TOTAL	100%

NET CONTENTS
32 Fl.Oz

EPA Registration Number: **97092-1**

EPA Est. No. **97092-CA-1**

KEEP OUT OF REACH OF CHILDREN

CAUTION:

SEE RIGHT SIDE FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND FIRST AID



Questions or Comments?

Please visit www.qyk.com or call 888-586-6386

Manufactured by: Qyk Brands LLC,
12101 Western Ave, Garden Grove CA 92841

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS Caution. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice. Have the container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information.

STORAGE AND DISPOSAL Store in original container in areas inaccessible to children. Nonrefillable container. Do not reuse or refill this container. Wrap and put in trash or offer for recycling if available.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

TO CLEAN AND DEODORIZE:

Spray directly on surface to be cleaned. Wipe or allow to air dry.

DISINFECTION / VIRUCIDAL* / FUNGICIDAL / MOLD & MILDEW CONTROL DIRECTIONS:

Spray 6 - 8 inches from surface. Avoid breathing spray. Treated surfaces must remain visibly wet for 10 minutes. For SARS-CoV-2, treated surfaces must remain visibly wet for 3 minutes. For Influenza Virus Type A and Human Coronavirus, treated surfaces must remain visibly wet for 1 minute. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a pre cleaning is required.

Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

For fungicidal activity: Dr J's Surface Disinfectant Spray is an effective one-step fungicide against Trichophyton mentagrophytes when used on hard, non-porous surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and

exercise facilities. Allow surface to remain visibly wet for 10 minutes.

For mold and mildew: Repeat treatment every 7 days or more often if new growth appears. This product is not for use on medical device surfaces.

***KILLS HIV-1 AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 [HIV-1] [associated with AIDS] or Hepatitis B Virus [HBV] or Hepatitis C Virus [HCV].

Special Instructions for Cleaning and Decontamination against HIV-1 or HBV or HCV on Surfaces / Objects Soiled With Blood / Body Fluids. Personal Protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection. Cleaning Procedure: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of [Dr.'s] [QYK] Surface Disinfectant [Spray]. Contact Time: Leave surface wet for 1 minute [60 seconds] for HIV-1 and 10 minutes for HBV and HCV. Use a 10 minute contact time for disinfection against all other viruses, bacteria and fungi claimed. Disposal of Infectious Material: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

Dr.J's SurfaceDisinfectant is a one-step Hospital use disinfectant⁶, Bactericidal tested according to the current AOAC Use-Dilution Test Method, Fungicidal according to the AOAC Fungicidal Test and Virucidal* tested according to the virucidal qualification modified in the presence of 5% organic serum against:

Bacteria: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Acinetobacter baumannii, Burkholderia cepacia Campylobacter jejuni, Corynebacterium ammoniagenes Enterobacter aerogenes Enterobacter cloacae NDM-1 Enterococcus faecalis, Enterococcus faecalis - Vancomycin resistant Escherichia coli, Escherichia coli, New Delhi Metallo-Beta Lactamase (NDM-1) ESBL Escherichia coli, Klebsiella pneumoniae, Klebsiella pneumoniae - NDM-1 positive Legionella pneumophila Listeria monocytogenes, Salmonella schottmuelleri, Salmonella typhi, Serratia marcescens Shigella dysenteriae, Staphylococcus aureus - Community Associated Methicillin-Resistant, Staphylococcus aureus - Community Associated Methicillin-Resistant, Staphylococcus aureus -

Methicillin-Resistant, Staphylococcus aureus Multi-Drug Resistant, Staphylococcus aureus - Vancomycin Intermediate Resistant - [VISA] Streptococcus pyogenes, Vibrio cholerae

Animal Viruses: Avian Infectious Bronchitis Virus Avian Influenza (H3N2), Avian Influenza (H5N1), Canine Distemper Virus, Feline Calicivirus, Murine Norovirus³, Newcastle's Disease Virus, Pseudorabies Virus.

Fungi: Aspergillus Niger⁴, Aspergillus fumigatus, Trichophyton mentagrophytes.

Viruses: *Hepatitis B Virus, *Hepatitis C Virus, *Herpes Simplex Virus Type 1, *Herpes Simplex Virus Type 2, *HIV-1, *Human Coronavirus *Influenza Virus Type A / Brazil, *Influenza Virus Type A / Hong Kong, *2013 Influenza A Virus (H7N9) *Respiratory Syncytial Virus,*Vaccinia, *Norwalk Virus *Rotavirus, *SARS Associated Coronavirus,

*SARS-Related Coronavirus 2 (SARS-CoV-2) cause of COVID-19

¹ SARS-Related Coronavirus 2

² On hard, non-porous, non-food surfaces

³ Tested in deionized water

⁴ follow mold/mildew directions

⁵ When used according to the disinfection directions.

⁶ When used according to the sanitization directions.

⁷ Staphylococcus aureus

⁸ Pseudomonas aeruginosa

⁹ Salmonella enterica

¹⁰ Allow the surface to adjust to room temperature before disinfection.



BARCODE

Batch Number:
Date Packaged: